# EVIDENCE FOR THE EVOLUTION OF YOUNG EARLY-TYPE GALAXIES IN THE GOODS/CDF-S FIELD

THOMAS H. PUZIA,<sup>1,2</sup> BAHRAM MOBASHER,<sup>2,3</sup> AND PAUL GOUDFROOIJ<sup>2</sup> Received 2006 September 25; accepted 2007 May 25

# ABSTRACT

We have developed an efficient photometric technique for identifying young early-type galaxy candidates using a combination of photometric redshifts, spectral-type classification, and optical/near-infrared colors. Applying our technique to the GOODS *HST* ACS and VLT ISAAC data, we have selected a complete and homogeneous sample of young elliptical candidates among early-type field galaxies. The distribution of structural parameters for these candidates shows that their selection, which is based on early spectral types, is fully consistent with early morphological types. We investigate the evolution of their luminosities and colors as a function of redshift and galaxy mass and find evidence for an increasing starburst mass fraction in these young early-type galaxy candidates at higher redshifts. Moreover, we find indications for a systematically larger young elliptical fraction among sub- $L^*/2$  early-type galaxies compared to their brighter counterparts. The total fraction among the field early-type galaxies increases with redshift, irrespective of galaxy luminosity. Our results are most consistent with galaxy formation scenarios in which stars in massive early-type field galaxies are assembled earlier than in their low-mass counterparts.

*Key words:* galaxies: elliptical and lenticular, cD — galaxies: evolution — galaxies: formation *Online material:* color figures

# 1. INTRODUCTION

The last star formation burst in galaxies defines their photometric appearance and is an important diagnostic of galaxy formation and evolution. The class of rejuvenated early-type galaxies includes so-called K+A galaxies,<sup>4</sup> which are objects that show spectroscopic signatures of old (K-type) and young (A-type) stellar populations (Dressler & Gunn 1983). The lack of emission lines in their spectra indicates that star formation processes abruptly ceased ~1 Gyr ago, followed by a quiescent evolution into normal early-type galaxies (e.g., Couch & Sharples 1987).

Several scenarios have been suggested to explain this observed rejuvenation phenomenon: (1) galaxy-galaxy mergers (e.g., Zabludoff et al. 1996), (2) interactions of infalling galaxies with the intracluster medium (e.g., Gunn & Gott 1972; Bothun & Dressler 1986), (3) galaxy harassment (e.g., Moore et al. 1996, 1998), (4) tidally induced star formation (e.g., Byrd & Valtonen 1990), and (5) dusty starbursts (e.g., Poggianti et al. 1999; Poggianti & Wu 2000). While tidal interaction and galaxy harassment are important events in the overall context of galaxy formation and evolution (e.g., Bekki et al. 2005), they do not provide an obvious explanation for the sudden halt of star formation without invoking additional processes. Ram pressure stripping and mergers are currently seen as the most viable scenarios to describe the ignition and sudden cessation of star formation in young early-type galaxies (e.g., Rose et al. 2001; Bekki & Couch 2003; Shioya et al. 2002, 2004; Quintero et al. 2004). Moreover, recent radio observations appear to exclude a dusty starburst scenario in general, as the measured radio fluxes are inconsistent with high star formation rates (i.e.,  $\gtrsim 10-100 \ M_{\odot} \ yr^{-1}$ ) for virtually all local poststarburst early-type galaxies (Miller & Owen 2001; Goto 2004). However, Smail et al. (1999) do find evidence for faint radio fluxes in spectroscopically confirmed poststarburst galaxies in distant dense clusters at  $z \approx 0.4$ .

K+A fractions are known to vary significantly with both redshift and environmental density. Several studies have revealed that the brightest K+A galaxies in nearby clusters are sub-L\* systems (Caldwell et al. 1999; Poggianti et al. 2004), while at intermediate redshifts ( $z \ge 0.8$ ) K+A galaxies are found with  $L \approx 3L^*$ (Dressler et al. 1999; Tran et al. 2003), suggesting an increase of K+A cluster galaxy mass with redshift. So far, the perhaps most unbiased comparison between the K+A fractions in field and cluster environments has been done by Tran et al. (2003, 2004), who found a field K+A fraction of  $\sim$ 3% and a factor  $\sim$ 4 higher fraction in cluster environment for the redshift range 0.3 < z < 1. Tran et al. further noted that the field K+A fraction shows strong fluctuations and is sensitive to selection criteria. Furthermore, the relatively short lifetime of the K+A signature (~1 Gyr; Couch & Sharples 1987; Belloni et al. 1995; Barger et al. 1996) implies that these galaxies are rare. This indicates the need for wide-area surveys to identify statistically significant samples of young earlytype galaxies and determine their number fractions in both field and cluster environments. If their selection is performed consistently between data sets and galaxy formation models, their abundance may be a sensitive diagnostic for star formation activity in field and cluster galaxies as a function of redshift, making these rejuvenated objects useful tools to constrain hierarchical galaxy formation scenarios.

Keeping in mind that medium-resolution spectroscopy of all galaxies in large-area surveys is very expensive in terms of telescope time, the aim of this paper is to develop a photometric technique to identify young early-type galaxy *candidates* (hereafter yE candidates) for spectroscopic follow up using a combination of photometric redshifts, spectral-type classification, and optical/nearinfrared color-color diagrams. Applying this technique to the

<sup>&</sup>lt;sup>1</sup> Herzberg Institute of Astrophysics, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada.

<sup>&</sup>lt;sup>2</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA.

<sup>&</sup>lt;sup>3</sup> Affiliated with the Space Sciences Department of the European Space Agency.

<sup>&</sup>lt;sup>4</sup> Historically, these galaxies were named "E+A" galaxies because of their early-type morphology appearance on the very first images. However, subsequent studies have shown that many of these galaxies contain significant disk components, so that the "E" becomes unjustified and is substituted by "K," owing to the spectral type of the underlying old stellar population (Franx 1993).

GOODS/CDF-S data set (Giavalisco et al. 2004), we select a complete and homogeneous sample of yE candidates in the redshift range  $0 < z \le 1$ . Finally, we investigate correlations of the colors of yE candidate galaxies with redshift and galaxy mass, which, if confirmed by spectroscopy, may put strong constraints on galaxy formation models. We use the standard  $\Lambda$ CDM cosmology parameters  $\Omega_M = 0.3$ ,  $\Omega_{\Lambda} = 0.7$ , and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$  throughout this work.

### 2. GOODS/CDF-S DATA

One of the two fields covered by the Great Observatories Origins Deep Survey (GOODS) Treasury Program is the Chandra Deep Field–South (CDF-S), with optical *HST* ACS and groundbased VLT ISAAC near-infrared photometry (Giavalisco et al. 2004). These data span the wavelength range from near-UV to the near-infrared and provide ACS imaging with very high spatial resolution (0.03") for ~86,000 galaxies down to  $V \approx 27.5$ . Object detection and photometry were performed with SExtractor (Bertin & Arnouts 1996) after the images in all passbands were convolved to a common FWHM  $\approx 0.45$ ". The final magnitudes are the MAG\_AUTO SExtractor magnitudes, which measure the total light.

Using the multi–wave band data available for galaxies in this field, photometric redshifts were measured following the technique presented in Mobasher et al. (2007). Briefly, this approach compares the template spectral energy distributions (SEDs) for different spectral types of galaxies, shifted in redshift space, with the observed SED for galaxies detected in GOODS (see also § 4.1);  $\chi^2$  fits are performed, and the redshift and spectral type corresponding to the minimum  $\chi^2$  value are associated with those for the galaxy in question. The template spectral types consist of elliptical, Sbc, Scd, and Im-type galaxies from Coleman et al. (1980) and starburst templates from Kinney et al. (1996). Bayesian priors based on observed galaxy luminosity functions are used (as detailed in Mobasher et al. 2007). Comparison with galaxies with available spectroscopic redshifts yields a photometric redshift accuracy of  $\sigma(\Delta z) = 0.03$ , where  $\Delta z = (z_{phot} - z_{spec})/(1 + z_{spec})$ .

In order to check the accuracy of our spectral types, we compare these with the morphology of galaxies measured from the GOODS-S ACS images. The morphologies are estimated using the concentration and asymmetry parameters (Conselice 1997; Abraham et al. 1996), measured for individual galaxies (B. Mobasher, 2007 private communication). We found a very good correlation between the morphology parameters and spectral types, particularly for elliptical galaxies. Therefore, we have reliable identification of elliptical galaxies, based on their spectral types, which constitutes the main selection criterion for the present study.

We use the information on redshifts and spectral types to compute *K*-corrections for all galaxies in our sample. Throughout the rest of this work we use AB magnitudes (Oke 1974)<sup>5</sup> and correct for Galactic foreground reddening  $E_{B-V} = 0.008$  derived from the Schlegel et al. (1998) maps. We select galaxies brighter than  $M_K = -19.0$  mag from the initial GOODS/CDF-S sample, which have a mean photometric redshift accuracy of  $\Delta z \le 0.15$ , corresponding to a 95% confidence interval. This photometric redshift uncertainty limit is introduced in order to keep the average color uncertainty below 0.15 mag due to *K*-corrections (see also § 3). The final sample is volume-selected and complete to  $z \approx 1$ . We exclude objects where the photometry was compromised by saturated pixels, or corrupted or truncated isophotal apertures, and where the extraction algorithm experienced problems during deblending or the photometric errors were larger than  $\Delta m = 0.15$  mag in any filter. The final catalog includes objects with *HST* ACS *BViz* and VLT ISAAC *JHKs* magnitudes.<sup>6</sup>

# 3. STELLAR POPULATION MODELS

#### 3.1. Diagnostic Colors

Based on the Bruzual & Charlot (2003, hereafter BC03) simple stellar population (SSP) models and their 2003 version of the GALAXEV stellar population synthesis code, we compute optical/ near-infrared photometric colors for composite stellar populations. We use the Salpeter IMF with limits at 0.1 and 100  $M_{\odot}$ . By comparison with the control sample of spectroscopically confirmed K+A galaxies, we identify the best diagnostic combination of optical and near-infrared colors to select yE candidates.

Given the availability of many filters in our GOODS data set (BVizJHK<sub>s</sub>), we investigate several filter combinations that may most efficiently reduce the age-metallicity-extinction degeneracy in the color-color plane. Due to the presence of a hotter  $\sim 1$  Gyr young stellar population, the B-band flux of yE galaxies is expected to be enhanced relative to older ( $\gg1$  Gyr) stellar populations. For slightly younger stellar populations with ages  $\sim 0.4$ -0.8 Gyr, the asymptotic giant branch (AGB) is densely populated with relatively cool stars (Renzini 1981), causing a stronger output in the near-infrared K band (Persson et al. 1983). Since a small population of luminous thermally pulsing AGB stars (TP-AGB) can significantly contribute to the integrated light of galaxies, the consideration of such short-lived evolutionary phases in SSP models is essential (Maraston 1998; Mouhcine & Lançon 2003). Past this so-called AGB phase transition, near-infrared light is primarily sensitive to the mean temperature of the red giant branch, which is mainly driven by the luminosity-weighted mean metallicity of composite stellar populations.

In Figure 1 we show the effective filter throughputs for HST ACS and the VLT ISAAC instrument together with SEDs for single-burst stellar populations for ages t = 290 Myr, 640 Myr, 1.4 Gyr, 2.5 Gyr, and 11 Gyr at solar metallicity. For illustration purposes all SEDs were normalized to the flux at 1.3  $\mu$ m. It is clear from the figure that the B band contains the most agesensitive flux and that the B - J color is a very good age indicator. It is instructive to note that the Lyman break is a sensitive indicator for identifying constantly star-forming or very young poststarburst galaxies (i.e.,  $t \leq 0.5$  Gyr; see the dotted SED in Fig. 1). However, the Balmer break is the more sensitive proxy for identifying poststarburst galaxies with intermediate ages (i.e.,  $t \ge 0.5$  Gyr), such as K+A galaxies. To find a good metallicity indicator we need to move to the near-infrared, past the z band, where the continuum slope shows little impact of varying age. Here, the H-band continuum suffers from variations due to strong water vapor absorption, which is sensitive to the mean temperature of the stellar population (i.e., age sensitivity). However, the relative continuum fluxes in J and Kshow little variation with age, and we define J - K as our best metallicity indicator, which exhibits only modest age sensitivity during the AGB phase transition (see also Ferraro et al. 2000).

The best choice of the most age-sensitive and the most metallicity-sensitive color also depends on data quality. For reference, the average color uncertainty introduced by the photometric redshift error is  $\leq 0.15$  mag in both B - J and J - K for all galaxy types at  $z \leq 1$ . Given the mean uncertainty of all color combinations of our data, the most age-sensitive color is B - J, and the most metallicity-sensitive color is J - K, which is virtually

<sup>&</sup>lt;sup>5</sup> AB magnitudes are defined by  $m_{AB} = -2.5 \log f_{\nu} - 48.6$ , where  $f_{\nu}$  is in ergs s<sup>-1</sup> cm<sup>-2</sup> Hz<sup>-1</sup>. Alternatively, one can write  $m_{AB} = -2.5 \log f_{\nu} + 8.9$ , where  $f_{\nu}$  is the flux in janskys.

<sup>&</sup>lt;sup>6</sup> In the following we refer to the  $K_s$  filter as the K band.



FIG. 1.—Comparison of rest-frame solar-metallicity SEDs for single-burst stellar populations with ages t = 290 Myr, 640 Myr, 1.4 Gyr, 2.5 Gyr, and 11 Gyr. A dotted line illustrates the SED of a solar-metallicity stellar population with a constant star formation rate after 6 Gyr. All SEDs were taken from BC03. Total filter throughputs for ACS and ISAAC, including mirror reflectivity, window transmission, and detector quantum efficiency, are shown at the top of the panel.

independent of dust reddening. This *BJK* combination optimizes the selection of poststarburst intermediate-age stellar populations at redshifts  $z \leq 1$ . For higher redshifts, the age-metallicity degeneracy dilutes the age/metallicity sensitivity of this color combination as the rest-frame flux at  $\lambda < 1 \ \mu$ m moves into the *J* band, and a different filter combination is required.

A comparable photometric selection technique, although specifically targeting star-forming galaxies at higher redshifts, was described by Daddi et al. (2004). Similar techniques to detect intermediate-age globular clusters (i.e., single-burst stellar populations) in nearby galaxies have recently been used by Goudfrooij et al. (2001), Puzia et al. (2002), and Hempel & Kissler-Patig (2004).

#### 3.2. Spectroscopically Confirmed K+A Galaxies

To test our photometric selection technique we use the K+A galaxy sample of Galaz (2000). This data set contains spectroscopically confirmed K+A galaxies in nearby ( $z \approx 0.05$ ) and distant ( $z \approx 0.3$ ) clusters and in the nearby field ( $z \approx 0.1$ ). All galaxies have been classified as K+A (Couch & Sharples 1987; Franx 1993; Caldwell & Rose 1997; Zabludoff et al. 1996), without any indications of emission lines. The near-infrared photometry of the Galaz study was performed with the Las Campanas NICMOS3 Hg:Cd:Te detector in the CIT filter system (Persson et al. 1998). We compute the transformations to the Bessell & Brett (1988, hereafter BB) standard system using BC03 SEDs with ages from 0.2 to 15 Gyr and [Fe/H] values from -2.25 to +0.4 dex, and find the following BB-CIT transformations:

$$J_{\rm BB} = J_{\rm CIT} - 0.0045 + 0.0102(J - K)_{\rm CIT},$$
  
$$(J - K)_{\rm BB} = 0.0147 + 0.9835(J - K)_{\rm CIT},$$

with an rms scatter of 0.0009 and 0.007 mag, respectively. Galaz's Table 2 lists Johnson *B* magnitudes as reported in the NED<sup>7</sup> at the time, likely coming from various original sources. We compare these *B* magnitudes with those in the Automatic Plate Measuring



FIG. 2.—Comparison of *B* magnitudes for spectroscopically confirmed K+A galaxies taken from Galaz (2000) with those taken from the APM catalog. The vertical solid line shows the mean formal uncertainty.

(APM) catalog<sup>8</sup> (Maddox et al. 1990) and show the comparison plot in Figure 2. We find significant offsets for the optical photometry between the Galaz and APM catalogs. Since the magnitudes in the APM catalog were derived in an internally consistent way, we adopt the APM *B* magnitudes for further analysis in this paper. The mean uncertainties in *BJK* colors of the comparison sample are 0.12, 0.10, and 0.18 mag, respectively. All optical and near-infrared magnitudes from the GOODS catalog and the comparison sample represent total magnitudes.

# 3.3. Confirmed K+A Galaxies in the BJK Color-Color Plane

In Figure 3 we compare  $(B - J)_{AB}$  versus  $(J - K)_{AB}$  rest-frame colors of spectroscopically confirmed K+A galaxies with SSP model predictions from two different groups: BC03 and Anders & Fritze-v. Alvensleben (2003, hereafter AF03). The BC03 predictions were calculated in the AB system with the same filter transmission functions as the GOODS/CDF-S data. For the AF03 models, which are tabulated in Vega-based magnitudes using the BB filter system, we follow the Vega-AB transformations:  $K_{AB} =$  $1.891 + K_{\text{Vega}}, J_{\text{AB}} = 0.91 + J_{\text{Vega}}, \text{ and } B_{\text{AB}} = -0.105 + B_{\text{Vega}}.$ As expected, the location of confirmed K+A galaxies in the colorcolor plane corresponds to old to intermediate ages. We find that the absolute age calibration differs between the two models. The BC03 models predict (luminosity-weighted) ages between  $\sim 15$ and  $\sim 2$  Gyr for most K+A galaxies, while the models of AF03 predict younger ages in the range  $\sim 1-5$  Gyr. The metallicity scales, on the other hand, are very similar in both models. The majority of confirmed K+A galaxies have luminosity-weighted metallicities between  $[Z/H] \approx -0.7$  and 0.0 dex, with a few outliers at supersolar metallicities, which might be reddened objects.

#### 3.4. Composite Stellar Populations in the BJK Color-Color Plane

Several groups compared the mean colors of K+A galaxies using optical and near-infrared photometry with population synthesis models (e.g., Newberry et al. 1990; Belloni et al. 1995; Barger et al. 1996; Shioya et al. 2002; Balogh et al. 2005). In this work we combine spectral-type fitting with model predictions and investigate the influence of different metallicities and mass fractions of the starburst population on colors and luminosities of poststarburst galaxies in the optical/near-infrared color-color plane and in color-magnitude diagrams.

<sup>&</sup>lt;sup>7</sup> The NASA/IPAC Extragalactic Database, http://nedwww.ipac.caltech.edu.

<sup>&</sup>lt;sup>8</sup> See http://www.ast.cam.ac.uk/~apmcat.



FIG. 3.—Left: Comparison of the  $(B - J)_{AB}$  and  $(J - K)_{AB}$  rest-frame colors of spectroscopically confirmed K+A galaxies (*open symbols*) with the SSP models of BC03. Squares mark cluster K+A galaxies, and circles show field K+A galaxies. Isochrones (*solid lines*) are plotted for ages t = 200, 500, and 700 Myr, and 1, 1.3, 2, 5, and 15 Gyr. Isometallicity lines (*dotted lines*) are shown for the metallicities [Z/H] = -2.3, -1.7, -0.7, -0.4, 0.0, and +0.4 dex. Right: Same data, but this time with the SSP models of AF03. Here isochrones (*solid lines*) were plotted for the ages t = 200, 500, and 700 Myr, and 1, 1.3, 2, 5, and if (Z/H] = -1.7, -0.7, -0.4, 0.0, and +0.4 dex. Each panel has the extinction vector for  $E_{B-V} = 0.2$  mag indicated in the upper left corner. The mean uncertainties of the Galaz (2000) data are shown in the upper right corner. [See the electronic edition of the Journal for a color version of this figure.]

Figure 4 shows the evolution of BC03 isochrones where a starburst is ignited after 10 Gyr of passive SSP evolution. Assuming a minor merger, we assign 10% of the final stellar mass to the starburst population. We adopt two metallicities, Z = 0.04 (twice solar metallicity) and  $Z = 10^{-4}$  ([Z/H] = -2.3 dex), for the starburst stellar population, which are superposed on top of the full range of metallicities ([Z/H] = -2.3 to +0.4 dex) of the underlying, older stellar population. Note the significant difference in B - J colors during the first ~1.5 Gyr after the starburst. If determined relative to passively evolving galaxies, this color offset is a sensitive poststarburst indicator; this is discussed below in more detail.

In Figure 4 (right) we show the influence of changing starburst mass fraction on the poststarburst galaxy colors. This time we choose solar metallicity for the starburst with mass fractions are 5%, 10% (minor merger), and 50% (major merger) of the final total stellar mass. The figure illustrates that the higher the starburst mass fraction the bluer the B - J color is during the poststarburst phase, in agreement with previous studies (e.g., Barger et al. 1996). In addition, there is a mass fraction-metallicity degeneracy of poststarburst near-infrared colors. Metal-rich major starbursts can exhibit the same J - K colors as metal-poor minor starbursts. This degeneracy is most prominent for intermediate metallicities in the range  $-0.7 \leq [Z/H] \leq 0$ . For more extreme metallicities the burst strength can be constrained with increasing confidence. In other words, yE galaxies with starburst mass fractions  $\leq 10\%$  and  $J - K \gtrsim 0$  are likely to have experienced a recent starburst with supersolar metallicity. The same mass-fraction limit and  $J - K \lesssim$ -0.3 mag indicate subsolar starburst metallicities.

In summary, B - J is mainly degenerate in age and starburst mass fraction, and J - K is mainly degenerate in metallicity and starburst mass fraction. We conclude that the B - J versus J - Kcolor-color plane can be used to identify poststarburst galaxies. But only extreme metallicities of the poststarburst stellar population can be narrowed down robustly. The color-color plane is not suitable to derive starburst mass fractions for intermediate metallicities without any other information. However, these degeneracies can be partly lifted by including rest-frame *K*-band luminosities. Our findings are relatively independent of the choice of SSP model, as the systematics in the AF03 models give the same results.

#### 4. SELECTION OF yE CANDIDATES

# 4.1. Spectral Types

In the following we develop a robust method to identify yE galaxy candidates from the GOODS/CDF-S photometric catalog using a combination of photometric redshift, spectral-type classification, and optical/near-infrared colors. The photometric redshifts and spectral types are measured by using template SEDs, shifting them in redshift steps and performing  $\chi^2$  fits to the observed SEDs at each step. The template corresponding to the minimum  $\chi^2$  value is selected, with its spectral type and redshift assigned to the galaxy in question. We use the observed templates for elliptical, Sbc, Scd, and Im-type galaxies from Coleman et al. (1980) and the starburst SEDs from Kinney et al. (1996). In order to increase the resolution in spectral type, we also divide the interval between adjacent spectral classes into two intermediate bins. For instance, between the two discrete spectral types for elliptical (T = 1) and Sbc-type spectra (T = 2), we include two discrete subcategories  $(T = 1\frac{1}{3}, 1\frac{2}{3})$  that are linear mixtures of the two boundary types. This allows a more continuous spectral classification across the Hubble types while not causing degeneracy. The choice of steps of  $\frac{1}{2}$  was made as a compromise between the goal to increase the resolution of spectral types on one hand and the goal to avoid creating degeneracies in the phot-z routine



FIG. 4.—Left: Evolution of  $(B - J)_{AB}$  vs.  $(J - K)_{AB}$  rest-frame colors for an SSP with a starburst that ignites after 10 Gyr of passive evolution and decays for another 5 Gyr. Two set of isochrones with ages t = 10.1, 10.2, 10.5, 11, 11.5, and 14.9 Gyr are shown for the metallicities [Z/H] = 0.4 (solid lines) and -2.3 dex (dot-dashed lines) of the starburst stellar population. The starburst strength amounts to 10% of the total final stellar mass. *Right*: Evolution of colors for different starburst strengths/masses for a solar-metallicity burst. Isochrones for ages t = 10.2, 10.5, 11.5, and 14.9 Gyr are plotted with different line types (solid, dashed, and dotted lines) indicating the dependence on starburst mass  $M_{burst} = 0.05, 0.1$ , and  $0.5 \times M_{total}$ . For reference we overplotted in both panels the SSP model grid from Fig. 3 [*left*]. All predictions were computed using the SSP models of BC03. Spectroscopically confirmed K+A galaxies are as in Fig. 3.

on the other. Details of the photometric redshift and spectral-type measurement are presented in Mobasher et al. (2004, 2007). We plot the distribution of the discrete spectral types in Figure 5 and define all objects that were assigned slightly earlier spectral types  $(T = 1\frac{1}{3})$  than passively evolving galaxies (T = 1) as yE candidates, i.e., objects which are bluer than normal ellipticals but redder than normal early-type spirals. Their location in Figure 5 is marked by a hatched histogram. Because later galaxy types can exhibit similar *BJK* colors as genuine yE galaxies, we use the spectral-type information to exclude late-type star-forming galaxies and to include only galaxy candidates with early spectral types in the following analysis. The distribution of photometric redshifts of all objects is shown in Figure 6. Most of our yE candidates have redshifts between  $z \approx 0.2$  and 0.8, with a few objects at slightly higher redshifts,  $z \approx 1.0$ .

# 4.2. Morphologies

To investigate the yE candidate morphologies we cross-correlate the GOODS/CDF-S sample with the structural parameter classification of Ravindranath et al. (2006). Based on BViz HST ACS images, the authors computed CAS parameters (concentration, asymmetry, and smoothness, defined in Conselice 2003) for all our sample sources. Because of the redshift distribution of our sources, each ACS filter probes a different wavelength and therefore a different morphological (stellar) component in a galaxy. To allow a comparison of structural parameters that is independent of redshift, we compute the redshift-corrected rest-frame B-band morphologies by linearly interpolating the CAS parameters between the pivot wavelengths of each ACS filter (see Sirianni et al. 2005) according to the redshift of each source. In Figure 7 we show the distribution of rest-frame B-band CAS parameters for all sample galaxies subdivided by spectral type, as in Figure 5. We find that the overall spectral-type classification of our yE



FIG. 5.—Distribution of spectral types *T* that were assigned by the photometric redshift routine for all objects in our selected GOODS/CDF-S sample (for details on the photometric redshift fitting routine we refer the reader to Mobasher et al. [2004, 2007]). The locations of individual spectral types are labeled. A subpopulation of several objects have a slightly earlier spectral type  $(T = 1\frac{1}{3})$  than passively evolving galaxies (T = 1). These are our yE candidates and are marked in the plot with a hatched bin. [See the electronic edition of the Journal for a color version of this figure.]



FIG. 6.—Distribution of photometric redshifts for all selected objects in our GOODS/CDF-S photometric data set (for details on the photometric redshift routine see Mobasher et al. 2004). The parameterization by spectral type is identical to the one in Fig. 5. [See the electronic edition of the Journal for a color version of this figure.]

candidates is consistent with early-type morphologies, which confirms the robustness of our procedure for selecting early-type poststarburst galaxies. In particular, the distribution of the CAS concentration parameter indicates that very few yE candidate galaxies with later type morphologies were selected by our photometric selection.

### 4.3. BJK Color-Color Diagram

Figure 8 (left) shows the GOODS/CDF-S data in the BJK colorcolor plane. Different symbols depict different spectral types as assigned to each object by the photometric redshift fitting routine. Most objects with SEDs consistent with passive early-type galaxies clump around rest-frame  $(B - J)_{AB} \approx 1.9$  and  $(J - K)_{AB} \approx$ 0. Objects with slightly later SED types (our yE candidates) scatter toward bluer B - J colors and have intermediate J - K colors that place their mean locus between spiral and star-forming galaxies. We find that the spectral-type selection is consistent with the BJK color selection and morphological classification. Moreover, most vE candidates are consistent with the previously computed composite stellar population models for poststarburst galaxies. Only one yE candidate at  $(B - J)_{AB} \approx 1.6$  and  $(J - J)_{AB} \approx 1.6$ K)<sub>AB</sub>  $\approx 0.35$  deviates significantly from the model predictions and might be a dusty starburst. For reference, we plot the control sample of spectroscopically confirmed K+A galaxies (open symbols).

To provide a comparison independent of SSP model predictions, we show in Figure 8 (*right*) nearby early-type (taken from Michard 2005) and late-type galaxies (taken from Pérez-González et al. 2003a) in the same color-color plane as the GOODS/CDF-S data. We find a good agreement between the early-type galaxies in GOODS/CDF-S and the local galaxy sample. However, the smaller photometric errors of the local galaxy sample allow a more clearly detected offset of the spectroscopically confirmed K+A galaxies toward bluer B - J colors (i.e., younger ages) compared to the local early-type galaxies, which line up in a tight sequence. We also notice that fewer spiral (*squares*) and Irr galaxies (*triangles*) lie outside of the SSP grid, which is again likely, partly due to smaller photometric errors of the nearby sample. We investigate whether this disagreement may be due to systematics in the photometric redshift determination, but find no differences in the error distributions between spiral and irregular galaxies, in particular in J - K, that appear to be primarily responsible for the scatter. A matching between the photometric redshifts of our sample and currently available spectroscopic redshifts from the literature (Le Fèvre et al. 2004) results in good agreement (see Table 1). Although we compare rest-frame colors in both samples, the galaxies in our GOODS/CDF-S data are more distant than the nearby comparison sample. Evolutionary factors, such as possibly enhanced dust fractions, may be responsible for the enhanced scatter in our data. Deep mid-infrared imaging and optical to infrared spectroscopy should be able to resolve this issue.

We define the population of yE galaxy candidates as those objects that have slightly later spectral types ( $T = 1\frac{1}{3}$ ; see Fig. 5) and BJK colors that are inconsistent with passively evolving (the reddest) early-type galaxies. The selected yE candidates are marked in Figure 8 by gray circles with central black dots. This definition might exclude some genuine K+A galaxies, since the control sample contains some spectroscopically confirmed K+A galaxies (in galaxy clusters) with BJK colors consistent with passively evolving early-type galaxies. Taking into account the average photometric error, this fraction is expected to be less than  $\sim 15\%$ . We point out that even with the color, spectral-type, and morphology selection, our yE candidate sample may be contaminated by early-type spiral galaxies that can mimic the morphologies of elliptical galaxies (e.g., when in a largely face-on configuration) and the SEDs of yE candidates because of their low but continuous star formation rates. Based on the local comparison sample we estimate this contamination fraction between  $\sim 10\%$  and 30%, depending on the exact color selection. Again, deep follow-up spectroscopy should be able to address this contamination fraction.

# 5. DISCUSSION

#### 5.1. Color-Magnitude Diagrams

Poststarburst galaxies tend to be brighter in the optical than their passive counterparts (e.g., Tran et al. 2003), which is likely due to their additional ~1 Gyr old stellar population. However, optical photometry is not able to provide strong constraints on how the stellar masses of these rejuvenated galaxies compare with those of passively evolving galaxies. The *K*-band stellar mass-tolight ratio changes by a factor of ~3 from ~1.5 to 13 Gyr (see also Drory et al. 2004), which is about an order of magnitude smaller a change than in the optical;  $M/L_K$  changes by about a factor of 1.5 between 500 Myr and 1.5 Gyr (the K+A phase). Hence, relative to optical colors an  $M_K$  selection defines the best approximation to a stellar-mass-selected data set.

In Figure 9 we show color-magnitude diagrams for all earlytype galaxies ( $T \le 1.5$ ) and the selected yE candidates, as well as the K+A control sample and nearby ellipticals. Overplotted are color-magnitude relations for early-type galaxies taken from the large galaxy survey of Mobasher et al. (1986), in good agreement with the GOODS data and the nearby elliptical galaxy sample. We find no significant difference between red-sequence galaxies and yE candidates in the near-infrared J - K versus  $M_K$  diagram, but we detect a significant offset in the B - J versus  $M_K$  diagram, where the yE candidates have significantly bluer colors at the same  $M_K$  (stellar mass) compared to the red sequence.

We show the influence of changing luminosity-weighted age  $(\Delta t)$ , metallicity  $(\Delta Z)$ , total galaxy mass  $(\Delta M_{gal})$ , and starburst mass fraction  $(\Delta M_*)$  as vectors in both color-magnitude diagrams;



Fig. 7.—Distribution of rest-frame *B*-band CAS parameters (concentration, asymmetry, and smoothness; see Conselice 2003) taken from Ravindranath et al. (2006) for all selected objects in our GOODS/CDF-S photometric data set parameterized by their spectral type as in Fig. 5. Note the strong separation of yE candidates in the concentration parameter (*top left*) from galaxies with later spectral types. [*See the electronic edition of the Journal for a color version of this figure.*]



FIG. 8.—*Left: BJK* color-color diagram (using dereddened rest-frame AB magnitudes) of galaxies in the GOODS/CDF-S field. The data symbols are parameterized by the galaxy spectral types, and are early-type (*filled circles*), spiral (*filled squares*), irregular (*filled triangles*), and starbursts (*stars*). The yE galaxy candidates are marked by small dark dots. Spectroscopically confirmed K+A galaxies taken from Galaz (2000) are indicated by open symbols, where open squares mark cluster K+A galaxies and open circles show field K+A galaxies. Average total photometric errors, including uncertainties of the photometry and the photometric redshift determination, are indicated in the upper right corner for yE candidates (*solid cross*), and the mean photometric uncertainties for confirmed K+A galaxies. Open circles show nearby ellipticals from Michard (2005), open squares and triangles show Sab and Scd spirals, and stars show local starburst galaxies, with the latter two taken from Pérez-González et al. (2003a). The same spectroscopically confirmed K+A sample as in the left panel is overplotted for comparison. The solid error bar shows average uncertainties for the elliptical sample, while the dotted cross is for the later type galaxies. SSP models in both panels are identical to those in Fig. 4 (*left*).

 $\Delta t$  and  $\Delta M_{gal}$ , and  $\Delta Z$  and  $\Delta M_*$  are highly degenerate in the J - K versus  $M_K$  diagram. In fact, the  $\Delta t$  and  $\Delta M_{gal}$  vectors are almost parallel to the red sequence (see Fig. 9, *top*), and therefore have negligible influence on the scatter in the J - K color, which is mainly driven by metallicity and the starburst mass fraction. The relatively small scatter in this diagram hints at a small scatter in metallicity and starburst mass fraction and/or is a signature of a correlation between these parameters. This degeneracy can be partly lifted with the B - J versus  $M_K$  diagram (see below). Four yE candidates fall significantly off the red sequence in the J - K versus  $M_K$  diagram. They are likely to have experienced significantly different starburst events than the rest of the sample. One possible explanation is that the metallicity of their starburst population is significantly lower.

The most influential parameters in the B - J versus  $M_K$  diagram are age and the starburst mass fraction (besides the total galaxy mass). Both vectors are highly inclined relative to the red sequence, which is a graphic explanation of the scatter in this plot. Metallicity

TABLE 1 Comparison between Photometric and Spectroscopic Redshifts

Subsample	$\langle \Delta z \rangle^{\rm a}$	$\sigma$	Ν
Е	$-0.109 \pm 0.004$	0.061	14
yE	$-0.119 \pm 0.012$	0.083	7
Sab	$0.028\pm0.004$	0.134	37
Scd	$-0.061\pm0.046$	0.185	4
Starburst	$0.224\pm0.000$		1

<sup>a</sup>  $\langle \Delta z \rangle = \langle z_{\text{phot}} - z_{\text{spec}} \rangle.$ 

plays a negligible role. We use the B - J versus  $M_K$  diagram to discuss the redshift evolution of our yE candidates in the next section.

#### 5.2. Starburst Mass Evolution

In the hierarchical merging scenario of galaxy formation, massive structures are expected to form on extended timescales from smaller subunits with a considerable fraction of stars forming relatively recently (e.g., Springel et al. 2005). If, in contrast, more massive structures form first, as seen in the monolithic collapse scenario, the majority of stars form at high redshifts ( $z \ge 2$ ) in intense starbursts (e.g., Larson 1975). The fraction of yE signatures among massive early-type galaxies is a measure of dissipative merging activity (recently often named "wet merging"), which is expected to be a function of redshift in the hierarchical merging picture. In contrast, no such evolution is expected in the early monolithic collapse scenario out to  $z \sim 2$ . This difference is expected to depend on environment and to be more pronounced in the field (Benson et al. 2002), which we probe here with the GOODS data set. The fraction of yE galaxies and the strength of the yE signatures among massive early-type field galaxies as a function of redshift therefore puts strong constraints on galaxy formation models.

To investigate the redshift evolution of our yE candidates, we subdivide the sample into four redshift bins between z = 0 and 1. For each galaxy we determine the B - J offset with respect to the red sequence and plot the residuals versus their absolute *K*-band luminosity in Figure 10. Our analysis is limited in redshift space by the completeness of our GOODS data in the *K* band. A 10  $\sigma$  point source detection in the VLT ISAAC data is feasible down to  $m_K(AB) = 25.1$  mag (Giavalisco et al. 2004), which translates



FIG. 9.—Optical/near-infrared color-magnitude diagrams of early-type galaxies (*filled circles*) and yE candidates (*dotted circles*). Large open symbols mark spectroscopically confirmed K+A galaxies (Galaz 2000) in the field (*circles*) and cluster environment (*squares*). Small open circles show nearby elliptical galaxies from Michard (2005), as in Fig. 8 (*right*). The nearby data were corrected for passive evolution, according to BC03 models, to the reference redshift z = 0.5 of our GOODS data, which includes only a correction of  $\Delta(B - J) = 0.2$  mag. Arrows indicate the relative changes in  $\Delta t$  between 500 Myr and 1.5 Gyr (*short arrow*) and 500 Myr and 13 Gyr (*long arrow*),  $\Delta Z$  between 0.2Z and solar metallicity,  $\Delta M_{gal}$  between  $10^{10}$  and  $10^{11} M_{\odot}$ , and  $\Delta M_*$  between 5% and 50% of the starburst mass fraction. Color-magnitude relations for early-type galaxies from the survey of Mobasher et al. (1986) are overplotted as dashed lines. Average photometric errors of yE candidates (*solid cross*) and spectroscopically confirmed K+A galaxies (*dotted cross*) are plotted in the upper right corner of each panel. [*See the electronic edition of the Journal for a color version of this figure*.]

into  $M_{K,AB} = -19.0$  mag at a redshift of z = 1 (indicated by a vertical line in Fig. 10). Hence, completeness is not an issue for the following analysis.

We observe that at any given  $M_K$  the B - J residuals are systematically larger for higher redshift galaxies. Spectroscopically confirmed K+A galaxies show a similar trend, but their redshift range is significantly smaller ( $z \le 0.12$ ) than that of our sample  $(z \le 1; \text{see Fig. 6})$ . Note that confirmed field K+A galaxies show larger  $\delta(B - J)$  than confirmed cluster K+A galaxies, which lie closer to the red sequence. We also find tentative evidence for a systematic difference in  $\langle \delta(J-K) \rangle$  between confirmed *field* K+A galaxies (open circles) and the combination of our vE candidates and confirmed *cluster* K+A galaxies. It appears that a metallicity difference is the most obvious way to explain this offset, given that in the more age-sensitive B - J panel the yE candidates lie among, or are bluer than, the nearby confirmed field and cluster K+A galaxies. However, the photometric errors of the spectroscopically confirmed K+A sample require that this intriguing result be confirmed with deep spectroscopy.

Parameters likely responsible for the evolution in  $\delta(B - J)$  with redshift are (1) a younger age of the poststarburst population at higher redshift or (2) a larger mass fraction of the poststarburst population at higher redshifts [we established in § 3 that metallicity has a smaller effect on  $\delta(B - J)$ ]. First we consider age effects. Under the assumption that the yE candidates indeed host 0.5-1.5 Gyr old poststarburst populations with strong Balmer absorption-line spectra, the vectors in Figure 10 show that the



Fig. 10.—Residual B - J colors with respect to the red sequence (see Fig. 9) as a function of absolute *K*-band magnitude for early-type galaxies. Dark dots mark yE candidates. Open symbols indicate spectroscopically confirmed K+A galaxies as in Fig. 9; taken from Galaz (2000). The symbol size is parameterized by redshift. All confirmed K+A galaxies are located at redshifts  $z \le 0.12$ . Vectors indicate the influence of varying age, metallicity, and starburst mass fraction, as in Fig. 9. Average photometric errors are indicated in the upper left corner of the panel, where the dotted cross indicates typical errors of the confirmed K+A galaxies and the solid cross shows errors for yE candidates. A long vertical line shows the 10  $\sigma$  point-source detection limit at z = 1. Small vertical lines indicate  $L^*$  at redshifts 0.2 and 1.0 for field galaxies (Drory et al. 2003). Note the paucity of yE candidates at magnitudes fainter than  $M_{K,AB} \approx -21.2$  and  $\delta(B - J)_{AB} \leq -0.8$  mag. [See the electronic edition of the Journal for a color version of this figure.]

difference in  $\delta(B - J)$  between yE candidates with  $z \leq 0.4$  and those with z > 0.6 is in principle similar to the effect of age fading from 0.5 to 1.5 Gyr. However, it is hard to imagine that poststarburst populations in yE galaxies at higher redshift (be it induced by galaxy interactions or by ram pressure stripping) would be systematically younger than in low-redshift ones, given the short duration of the vE phenomenon relative to the difference in look-back times within this redshift range. In the adopted WMAP cosmology, the difference in look-back times between z = 0.2and 1.0 is 7.8 - 2.4 = 5.4 Gyr, significantly longer than the duration of the yE phenomenon. Instead, one would expect the ages of such poststarburst populations to be randomly distributed (between 0.5 and 1.5 Gyr) at any redshift. We therefore suggest that it is more likely that the increase of yE candidates'  $\delta(B-J)$ with redshift is primarily due to an increasing mass fraction of the poststarburst population. If indeed the yE phenomenon identifies an important era in the assembly history of early-type galaxies, the lack of low-luminosity ( $M_K \gtrsim -21$ ) yE candidates at z > 0.6 in our GOODS sample would constitute strong evidence in favor of galaxy formation scenarios in which more massive early-type galaxies in the field are assembled earlier than their low-mass counterparts, which is in line with the "downsizing" picture (e.g., Cowie et al. 1996).

On the other hand, we currently cannot exclude the possibility that the higher redshift yE candidates do not actually host a (one) poststarburst population, but instead have systematically younger overall (luminosity-weighted) ages than the lower redshift yE candidates. More frequent bursts at higher redshifts, due to -

TABLE 2   Number Counts of E Galaxies and yE Candidates										
	SUB- <i>L</i> */2		SUPER- $L^*/2$							
Redshift	$N_{\rm E}$	$N_{\rm yE}$	$N_{\rm E}$	$N_{\rm yE}$	$f_{< L^*/2}{}^{\rm a}$	$f_{>L^*/2}^{\mathbf{b}}$				
>0.6	2	12	7	2	0.86	0.22				
< 0.6	18	19	16	2	0.51	0.11				

<sup>a</sup> Fraction  $f_{<L^*/2}$  of yE galaxy candidates among all sub- $L^*/2$  early-type

galaxies. <sup>b</sup> Fraction  $f_{>L^*/2}$  of yE galaxy candidates among all super- $L^*/2$  early-type galaxies.

more frequent mergers, might be a possible explanation, which would lead to systematically younger observed ages. Given the relevance of this issue in terms of its power to constrain early-type galaxy formation scenarios, we suggest a spectroscopic follow-up to test whether our yE candidates really do contain K+A features in their spectra.

### 5.3. Evolution of the yE Fraction

The fraction of poststarburst galaxies in nearby clusters was reported to increase with decreasing galaxy luminosity (e.g., Poggianti et al. 2001; Smail et al. 2001). The morphology bias prevents a direct comparison between the yE fractions in the cluster and field environment. We therefore select only early-type galaxies from our sample and investigate the yE fraction in the field. A similar trend of an increasing poststarburst fraction with decreasing galaxy luminosity is present in our GOODS sample of field galaxies. We quantify this trend by splitting the data at  $L^*/2$  along the pure luminosity evolution vector and counting the number of passive ellipticals and yE candidates. The corresponding values for the division for each redshift bin into super- $L^*/2$  and sub- $L^*/2$  systems are  $M_K^* = -22.27, -22.43, -22.64,$ and -22.75 mag for z = 0.3, 0.5, 0.7, and 0.9, respectively. Note that the samples are complete down to  $M_B = -16.3$  mag and  $M_K = -19.0$  mag at z = 1, which leaves our sample sensitive to objects with rest-frame colors  $B - K \leq 2.7$  mag but excludes extremely reddened objects, such as extremely red objects (EROs; e.g., Daddi et al. 2000). However, these galaxies are most frequent beyond redshift of unity (e.g., Georgakakis et al. 2006; Simpson et al. 2006) and are expected not to be a significant part of our initial sample. To improve sample statistics we merge the number counts of the two upper and two lower redshift bins, which results in a high- and low-redshift sample divided at z = 0.6. Galaxy number counts are given in Table 2.

We confirm the trend of a higher fraction of poststarbursts among early-type galaxies toward less luminous objects. In the highredshift sample we find a yE candidate fraction of 86% among all sub- $L^*/2$  early-type galaxies, whereas only 22% of the more luminous (massive) objects are yE candidates. The yE fractions shrink toward lower redshifts to 51% for sub- $L^*/2$  systems and 11% for their brighter counterparts. We suggest that these high yE fractions be confirmed (or refuted) using follow-up spectroscopy.

### 6. CONCLUSIONS

Based on the combination of photometric redshift, spectral-type classification, and optical/near-infrared colors, we have selected a sample of young early-type galaxy candidates (yE candidates) from the GOODS/CDF-S data set. Our technique relies on spectraltype fitting and color-color/color-magnitude diagnostic diagrams and provides an efficient method for selecting vE candidates using only photometric information. An analysis of CAS parameters shows that the selected yE candidates have early morphological types, which confirms that spectral-type and morphological-type selection are fully consistent with each other. The comparison of population synthesis models and observed properties of a sample of spectroscopically confirmed K+A galaxies provides strong circumstantial evidence that the selected candidates are genuine field early-type poststarburst galaxies.

We study the systematics that drive colors and magnitudes of poststarburst galaxies in diagnostic diagrams using current population synthesis models. Our analysis reveals evidence for a changing starburst mass fraction with increasing redshift in the sense that more massive/intense starbursts may be responsible for the yE signatures of massive field early-type galaxies at higher redshifts. Furthermore, we find a higher yE candidate fraction in sub- $L^*/2$ early-type galaxies compared to their more luminous counterparts. Within the redshift range of our data ( $z \leq 1$ ), we may be witnessing evidence for enhanced merging in the field toward higher redshifts, especially for high-luminosity galaxies. Similar results are obtained by studies of luminosity-weighted mean ages of nearby early-type galaxies, which indicate that low-L galaxies show a much greater scatter to younger luminosity-weighted mean ages than high-L galaxies (e.g., Caldwell et al. 2003; Pérez-González et al. 2003b; Sánchez-Blázquez et al. 2006). If follow-up spectroscopy of the yE candidates identified in this paper confirms their K+A nature, this suggests that the assembly of stars in high-L early-type field galaxies occurred earlier than in their lower L counterparts.

We thank Peter M. Pessev for his help in providing relevant near-infrared filter passband information. We are grateful to the referee, James Rose, for his insightful comments that improved the content of this paper. T. H. P. gratefully acknowledges support in the form of an ESA research fellowship and partial financial support through grant GO-10129 from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

#### REFERENCES

- Abraham, R. G., van den Bergh, S., Glazebrook, K., Ellis, R. S., Santiago, B. X., Surma, P., & Griffiths, R. E. 1996, ApJS, 107, 1
- Anders, P., & Fritze-v. Alvensleben, U. 2003, A&A, 401, 1063 (AF03)
- Balogh, M. L., Miller, C., Nichol, R., Zabludoff, A., & Goto, T. 2005, MNRAS, 360. 587
- Barger, A. J., Aragon-Salamanca, A., Ellis, R. S., Couch, W. J., Smail, I., & Sharples, R. M. 1996, MNRAS, 279, 1
- Bekki, K., & Couch, W. J. 2003, ApJ, 596, L13
- Bekki, K., Couch, W. J., Shioya, Y., & Vazdekis, A. 2005, MNRAS, 359, 949
- Belloni, P., Bruzual A., G., Thimm, G. J., & Roser, H.-J. 1995, A&A, 297, 61
- Benson, A. J., Ellis, R. S., & Menanteau, F. 2002, MNRAS, 336, 564

Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393

- Bessell, M. S., & Brett, J. M. 1988, PASP, 100, 1134 (BB)
- Bothun, G. D., & Dressler, A. 1986, ApJ, 301, 57
- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000 (BC03)
- Byrd, G., & Valtonen, M. 1990, ApJ, 350, 89
- Caldwell, N., & Rose, J. A. 1997, AJ, 113, 492
- Caldwell, N., Rose, J. A., & Concannon, K. D. 2003, AJ, 125, 2891
- Caldwell, N., Rose, J. A., & Dendy, K. 1999, AJ, 117, 140
- Coleman, G. D., Wu, C.-C., & Weedman, D. W. 1980, ApJS, 43, 393
- Conselice, C. J. 1997, PASP, 109, 1251
- . 2003, ApJS, 147, 1

- Couch, W. J., & Sharples, R. M. 1987, MNRAS, 229, 423
- Cowie, L. L., Songaila, A., Hu, E. M., & Cohen, J. G. 1996, AJ, 112, 839
- Daddi, E., Cimatti, A., & Renzini, A. 2000, A&A, 362, L45
- Daddi, E., Cimatti, A., Renzini, A., Fontana, A., Mignoli, M., Pozzetti, L., Tozzi, P., & Zamorani, G. 2004, ApJ, 617, 746
- Dressler, A., & Gunn, J. E. 1983, ApJ, 270, 7
- Dressler, A., Smail, I., Poggianti, B. M., Butcher, H., Couch, W. J., Ellis, R. S., & Oemler, A. J. 1999, ApJS, 122, 51
- Drory, N., Bender, R., Feulner, G., Hopp, U., Maraston, C., Snigula, J., & Hill, G. J. 2003, ApJ, 595, 698
- ——. 2004, ApJ, 608, 742
- Ferraro, F. R., Montegriffo, P., Origlia, L., & Fusi Pecci, F. 2000, AJ, 119, 1282
- Franx, M. 1993, ApJ, 407, L5
- Galaz, G. 2000, AJ, 119, 2118
- Georgakakis, A., Hopkins, A. M., Afonso, J., Sullivan, M., Mobasher, B., & Cram, L. E. 2006, MNRAS, 367, 331
- Giavalisco, M., et al. 2004, ApJ, 600, L93
- Goto, T. 2004, A&A, 427, 125
- Goudfrooij, P., Alonso, M. V., Maraston, C., & Minniti, D. 2001, MNRAS, 328, 237
- Gunn, J. E., & Gott, J. R. I. 1972, ApJ, 176, 1
- Hempel, M., & Kissler-Patig, M. 2004, A&A, 428, 459
- Kinney, A. L., Calzetti, D., Bohlin, R. C., McQuade, K., Storchi-Bergmann, T., & Schmitt, H. R. 1996, ApJ, 467, 38
- Larson, R. B. 1975, MNRAS, 173, 671
- Le Fèvre, O., et al. 2004, A&A, 428, 1043
- Maddox, S. J., Efstathiou, G., Sutherland, W. J., & Loveday, J. 1990, MNRAS, 243, 692
- Maraston, C. 1998, MNRAS, 300, 872
- Michard, R. 2005, A&A, 441, 451
- Miller, N. A., & Owen, F. N. 2001, ApJ, 554, L25
- Mobasher, B., Ellis, R. S., & Sharples, R. M. 1986, MNRAS, 223, 11
- Mobasher, B., et al. 2004, ApJ, 600, L167
- ------. 2007, ApJS, in press (astro-ph/0612344)
- Moore, B., Katz, N., Lake, G., Dressler, A., & Oemler, A. 1996, Nature, 379, 613
- Moore, B., Lake, G., & Katz, N. 1998, ApJ, 495, 139
- Mouhcine, M., & Lançon, A. 2003, A&A, 402, 425
- Newberry, M. V., Boroson, T. A., & Kirshner, R. P. 1990, ApJ, 350, 585
- Oke, J. B. 1974, ApJS, 27, 21

- Pérez-González, P. G., Gil de Paz, A., Zamorano, J., Gallego, J., Alonso-Herrero, A., & Aragón-Salamanca, A. 2003a, MNRAS, 338, 508
  - —. 2003b, MNRAS, 338, 525
- Persson, S. E., Aaronson, M., Cohen, J. G., Frogel, J. A., & Matthews, K. 1983, ApJ, 266, 105
- Persson, S. E., Murphy, D. C., Krzeminski, W., Roth, M., & Rieke, M. J. 1998, AJ, 116, 2475
- Poggianti, B. M., Bridges, T. J., Komiyama, Y., Yagi, M., Carter, D., Mobasher, B., Okamura, S., & Kashikawa, N. 2004, ApJ, 601, 197
- Poggianti, B. M., Smail, I., Dressler, A., Couch, W. J., Barger, A. J., Butcher, H., Ellis, R. S., & Oemler, A. J. 1999, ApJ, 518, 576
- Poggianti, B. M., & Wu, H. 2000, ApJ, 529, 157
- Poggianti, B. M., et al. 2001, ApJ, 563, 118
- Puzia, T. H., Zepf, S. E., Kissler-Patig, M., Hilker, M., Minniti, D., & Goudfrooij, P. 2002, A&A, 391, 453
- Quintero, A. D., et al. 2004, ApJ, 602, 190
- Ravindranath, S., et al. 2006, ApJ, 652, 963
- Renzini, A. 1981, Ann. Phys., 6, 87
- Rose, J. A., Gaba, A. E., Caldwell, N., & Chaboyer, B. 2001, AJ, 121, 793
- Sánchez-Blázquez, P., Gorgas, J., Cardiel, N., & González, J. J. 2006, A&A, 457, 809
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Shioya, Y., Bekki, K., & Couch, W. J. 2004, ApJ, 601, 654
- Shioya, Y., Bekki, K., Couch, W. J., & De Propris, R. 2002, ApJ, 565, 223
- Simpson, C., et al. 2006, MNRAS, 373, L21
- Sirianni, M., et al. 2005, PASP, 117, 1049
- Smail, I., Kuntschner, H., Kodama, T., Smith, G. P., Packham, C., Fruchter, A. S., & Hook, R. N. 2001, MNRAS, 323, 839
- Smail, I., Morrison, G., Gray, M. E., Owen, F. N., Ivison, R. J., Kneib, J.-P., & Ellis, R. S. 1999, ApJ, 525, 609
- Springel, V., et al. 2005, Nature, 435, 629
- Tran, K. H., Franx, M., Illingworth, G., Kelson, D. D., & van Dokkum, P. 2003, ApJ, 599, 865
- Tran, K. H., Franx, M., Illingworth, G. D., van Dokkum, P., Kelson, D. D., & Magee, D. 2004, ApJ, 609, 683
- Zabludoff, A. I., Zaritsky, D., Lin, H., Tucker, D., Hashimoto, Y., Shectman, S. A., Oemler, A., & Kirshner, R. P. 1996, ApJ, 466, 104